

- On *Plectranthus longitubus* MIQ. (*Aki-chôji*). Prov. Buzen : Mt. Ehiko (Sept. 3, 1934, E. TOBINAGA).
- 43) **Coleosporium Petasitis** LÉVEILLÉ in Ann. Sci. Nat. 3. Sér. VIII, p. 373, 1847.
On *Petasites japonicus* MIQ. (*Fuki*). Prov. Buzen : Mt. Ehiko (Sept. 3, 1934, E. TOBINAGA).
- 44) **Aecidium Hamamelidis** DIETEL in ENGL. Bot. Jahrb. XXVII, p. 571, 1899.
On *Hamamelis japonica* SIEB. et ZUCC. (*Mansaku*). Prov. Bungo : Mt. Kujyû (Aug. 14, 1931, O. ISHIUCHI).
- 45) **Aecidium Machili** P. HENNINGS in Hedwigia, XLI, p. (21), 1902.
On *Machilus Thunbergii* SIEB. et ZUCC. (*Tabunoki*). Prov. Chikuzen : Yamaguchi-mura (Aug. 27, 1931, O. ISHIUCHI).
- 46) **Aecidium Mori** BARCLAY in Jour. Asiatic Soc. Bengal, LX, pt. 2, p. 226, 1891.
On *Morus alba* L. (*Kuwa*) (*cult.*). Prov. Higo : Tochinoki (July 4, 1932, Y. HASHIOKA).
- 47) **Aecidium Paederiae** DIETEL in Hedwigia, XXXVI, p. 297, 1897.
On *Paederia chinensis* HANCE. (*Hekusokazura*). Prov. Chikuzen : Yamaguchi-mura (Aug. 29, 1931, O. ISHIUCHI).

Ixeris musashensis MAKINO and HISAUCHI, a new species

By

Kiyotaka HISAUCHI

久内清孝： にな属ニ一新種ヲ加フ

***Ixeris musashensis* MAKINO et HISAUCHI, nov. sp.**

Syn. *Lactuca musashensis* MAKINO, MSS.

Planta perennis tota glabra glauca. Caulis brevissimus ramosus pluri-
foliatus. Folia basilaria spatulata, oblanceolata vel lineari-oblonga subin-
tegra vel lacerata 3-8 cm. longa ca. 1 cm. lata basi cuneato-decurrentia valde
elongata apice rotundata vel acuta. Scapi 4-11 cm. alti. Bractea minima
ovata attenuata integra sed basi 1-2 denticulata. Capitula 1-2, 1½ cm.

lata. Caliculi 5 minimi ovati margine saepe dentati. Squamae involucri 7, lineari-lanceolatae apice acutae intus apice puberulae. Ligulae dilute lilacinae apice dentatae. Antherae atratae. Pappi albi setigeri, setis minutissimis erecto-patentibus.

Nom. Jap. *Takasago-nigana*.

Hab. NIPPON: Hondo—In graminosis circa Bushi in provincia Musashi. (6 V, 1922—Typus in Herb. Univ. Imp. Tokyo)

Planta endemica.

Present species is a small weed resembling *Ixeris chinensis* (THUNBERG) NAKAI⁽¹⁾ in its general appearance, particularly its colour of leaves and rays except that the scapes are aphyllous and much contracted as in *Ixeris stolonifer* A. GRAY,⁽²⁾ but has no runner. Found in the grassy region near Tokyo where *Ixeris chinensis* NAKAI, *I. japonica* NAKAI,⁽³⁾ *I. stolonifer* A. GRAY and *I. dentata* NAKAI⁽⁴⁾ were met with.

相當 = 古イ話デハアルガ 1922 年 (大正 11 年) 5 月 6 日牧野先生ト武州佛^{フツサ}子驛^シヲ徑テ青梅鐵道線ノ一驛福生ニ出タ事ガアツタ、其時たかさごさうダカにがなダカ恐ラク其間種ラシイモノガアツタノデ牧野先生ガ之ニタかさごにがなナル和名ヲ下シタ。其後武蔵野原野ノ各地ヲ稼行中時々之ニ逢着スルノデ其度毎ニ昔ノ事ヲ偲ンダノデアツタガ、喉元過グルト何ントカデ今迄忘レテ居タ處近頃余ノ標本箱ノ中カラ夫レガ出テ來タ。元來コノ植物ハ未ダ公ケニサレタモノデ無イカラ、無籍者デアル。然シ現在ノ牧野先生ハ新種ノ一ツ位見向キモサレナイシ、マタソソナ小乗のナ事ハ面倒ダト言フ様ナ風ナノデアルカラ止ムナク當時ノ同行者ト云フ資格デ之ヲ公ケニスルコトニシタ、其形狀ハ寫眞デ御覽ヲ願フトシテ置クガ尙一言スレバ舌狀花及花莖、葉等ノ色ハたかさごさうニ似テ居ルノミナラズ葉緣ノ形狀マデモ似テ居ルガ花莖ハにがなノ夫レヲ縮少シタカ又ハいはにがなノ夫レノ様デ細クテ短ク無葉デアル。何レににがな屬ノ某々間ノ自然間種デアラウガ、今其兩親ヲ引合ニ出スコトハ憶測ダカラ止メル。尤モ和名たかさごにがなハ當時ノ所感ニ依ツタモノデ臺灣ヲ聯想シタノデハナイ。

(1) *Ixeris chinensis* (TH.) NAKAI in Tokyo Bot. Mag. XXXIV, p. 152 (1920)

(2) *Ixeris stolonifer* A. GRAY, On the Botany of Japan in Memoirs Amer. Academy of Art and Science VI, p. 396 (1858-9)

(3) *Ixeris japonica* (TH.) NAKAI in Tokyo Bot. Mag. XL p. 575 (1926)

(4) *Ixeris dentata* (TH.) NAKAI in Flora Sylvatica Koreana Pars XIV, p. 114 (1923)

サテ春カラ秋マデ野外デ多數ノ同業者ノ姿ヲ見受ケルガ彼等ノ多クハ此寫眞ヲ見タラ肯カレル事デアラウ。尙原標本ハ一個シカ無イカラ東大腊葉室ニオク。



たかさごにがな *Ixeris musashensis* MAKINO et HISAUCHI. $\times 1/1$.